

# INVERSION IN THE PLANE BERKELEY MATH CIRCLE

ZVEZDELINA STANKOVA  
MILLS COLLEGE/UC BERKELEY  
SEPTEMBER 12 2006

## DEFINITION OF INVERSION IN THE PLANE

**Definition 1.** Let  $k(O, r)$  be a circle with center  $O$  and radius  $r$ . Consider a function on the plane,  $I : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ , sending a point  $X \neq O$  to the point on the half line  $OX^\rightarrow$ ,  $X_1$ , defined by

$$OX \cdot OX_1 = r^2.$$

Such a function  $I$  is called an *inversion of the plane* with center  $O$  and radius  $r$  (write  $I(O, r)$ .)

It is immediate that  $I$  is *not* defined at p.  $O$ . But if we compactify  $\mathbb{R}^2$  to a sphere by adding one extra point  $O_\infty$ , we could define  $I(O) = O_\infty$  and  $I(O_\infty) = O$ .

An inversion of the plane can be equivalently described as follows. If  $X \in k$ , then  $I(X) = X$ . If  $X$  lies outside  $k$ , draw a tangent from  $X$  to  $k$  and let  $X_2$  be the point of tangency. Drop a perpendicular  $X_2X_1$  towards the segment  $OX$  with  $X_1 \in OX$ , and set  $I(X) = X_1$ . The case when  $X$  is inside  $k$ ,  $X \neq O$ , is treated in a reverse manner: erect a perpendicular  $XX_2$  to  $OX$ , with  $X_2 \in k$ , draw the tangent to  $k$  at point  $X_2$  and let  $X_1$  be the intersection of this tangent with the line  $OX$ ; we set  $I(X) = X_1$ .

## PROPERTIES OF INVERSION

Some of the basic properties of a plane inversion  $I(O, r)$  are summarized below:

- $I^2$  is the identity on the plane.
- If  $A \neq B$ , and  $I(A) = A_1, I(B) = B_1$ , then  $\triangle OAB \sim \triangle OB_1A_1$ . Consequently,

$$A_1B_1 = \frac{AB \cdot r^2}{OA \cdot OB}.$$

- If  $l$  is a line with  $O \in l$ , then  $I(l) = l$ .
- If  $l$  is a line with  $O \notin l$ , then  $I(l)$  is a circle  $k_1$  with diameter  $OM_1$ , where  $M_1 = I(M)$  for the orthogonal projection  $M$  of  $O$  onto  $l$ .
- If  $k_1$  is a circle through  $O$ , then  $I(k_1)$  is a line  $l$ : reverse the previous construction.
- If  $k_1(O_1, r_1)$  is a circle not passing through  $O$ , then  $I(k_1)$  is a circle  $k_2$  defined as follows: let  $A$  and  $B$  be the points of intersection of the line  $OO_1$  with  $k_1$ , and let  $A_1 = I(A)$  and  $B_1 = I(B)$ ; then  $k_2$  is the circle with diameter  $A_1B_1$ . Note that the center  $O_1$  of  $k_1$  does *not* map to the center  $O_2$  of  $k_2$ .

Note that two circles are perpendicular if their tangents at a point of intersection are perpendicular; following the same rule, a line and a circle will be perpendicular if the line passes through the center of the circle. In general, the angle between a line and a circle is the angle between the line and the tangent to the circle at a point of intersection with the line.

- Inversion preserves angles between figures: let  $F_1$  and  $F_2$  be two figures (lines, circles); then

$$\angle(F_1, F_2) = \angle(I(F_1), I(F_2)).$$

## PROBLEMS

- (1) Given a point  $A$  and two circles  $k_1$  and  $k_2$ , construct a third circle  $k_3$  so that  $k_3$  passes through  $A$  and is tangent to  $k_1$  and  $k_2$ .
- (2) Given two points  $A$  and  $B$  and a circle  $k_1$ , construct another circle  $k_2$  so that  $k_2$  passes through  $A$  and  $B$  and is tangent to  $k_1$ .

- (3) Given circles  $k_1, k_2$  and  $k_3$ , construct another circle  $k$  which tangent to all three of them.
- (4) Let  $k$  be a circle, and let  $A$  and  $B$  be points on  $k$ . Let  $s$  and  $q$  be any two circles tangent to  $k$  at  $A$  and  $B$ , respectively, and tangent to each other at  $M$ . Find the set traversed by the point  $M$  as  $s$  and  $q$  move in the plane and still satisfy the above conditions.
- (5) Circles  $k_1, k_2, k_3$  and  $k_4$  are positioned in such a way that  $k_1$  is tangent to  $k_2$  at point  $A$ ,  $k_2$  is tangent to  $k_3$  at point  $B$ ,  $k_3$  is tangent to  $k_4$  at point  $C$ , and  $k_4$  is tangent to  $k_1$  at point  $D$ . Show that  $A, B, C$  and  $D$  are either collinear or concyclic.
- (6) Circles  $k_1, k_2, k_3$  and  $k_4$  intersect cyclicly pairwise in points  $\{A_1, A_2\}$ ,  $\{B_1, B_2\}$ ,  $\{C_1, C_2\}$ , and  $\{D_1, D_2\}$ . ( $k_1$  and  $k_2$  intersect in  $A_1$  and  $A_2$ ,  $k_2$  and  $k_3$  intersect in  $B_1$  and  $B_2$ , etc.)
- (a) Prove that if  $A_1, B_1, C_1, D_1$  are collinear (concylic), then  $A_2, B_2, C_2, D_2$  are also collinear (concylic).
- (b) Prove that if  $A_1, A_2, C_1, C_2$  are concyclic, then  $B_1, B_2, D_1, D_2$  are also concyclic.
- (7) (Ptolemy's Theorem) Let  $ABCD$  be inscribed in a circle  $k$ . Prove that the sum of the products of the opposite sides equals the product of the diagonals of  $ABCD$ :

$$AB \cdot DC + AD \cdot BC = AC \cdot BD.$$

Further, prove that for any four points  $A, B, C, D$ :  $AB \cdot DC + AD \cdot BC \geq AC \cdot BD$ . When is equality achieved?

- (8) Let  $k_1$  and  $k_2$  be two circles, and let  $P$  be a point. Construct a circle  $k_0$  through  $P$  so that  $\angle(k_1, k_0) = \alpha$  and  $\angle(k_2, k_0) = \beta$  for some given angles  $\alpha, \beta \in [0, \pi)$ .
- (9) Given three angles  $\alpha_1, \alpha_2, \alpha_3 \in [0, \pi)$  and three circles  $k_1, k_2, k_3$ , two of which do not intersect, construct a fourth circle  $k$  so that  $\angle(k, k_i) = \alpha_i$  for  $i = 1, 2, 3$ .
- (10) Construct a circle  $k^*$  so that it goes through a given point  $P$ , touches a given line  $l$ , and intersects a given circle  $k$  at a right angle.
- (11) Construct a circle  $k$  which goes through a point  $P$ , and intersects given circles  $k_1$  and  $k_2$  at angles  $45^\circ$  and  $60^\circ$ , respectively.
- (12) Let  $ABCD$  and  $A_1B_1C_1D_1$  be two squares oriented in the same direction. Prove that  $AA_1, BB_1$  and  $CC_1$  are concurrent if  $D \equiv D_1$ .
- (13) Let  $ABCD$  be a quadrilateral, and let  $k_1, k_2$ , and  $k_3$  be the circles circumscribed around  $\triangle DAC$ ,  $\triangle DCB$ , and  $\triangle DBA$ , respectively. Prove that if  $AB \cdot CD = AD \cdot BC$ , then  $k_2$  and  $k_3$  intersect  $k_1$  at the same angle.
- (14) In the quadrilateral  $ABCD$ , set  $\angle A + \angle C = \beta$ .
- (a) If  $\beta = 90^\circ$ , prove that that  $(AB \cdot CD)^2 + (BC \cdot AD)^2 = (AC \cdot BD)^2$ .
- (b) If  $\beta = 60^\circ$ , prove that  $(AB \cdot CD)^2 + (BC \cdot AD)^2 = (AC \cdot BD)^2 + AB \cdot BC \cdot CD \cdot DA$ .
- (15) Let  $k_1$  and  $k_2$  be two circles intersecting at  $A$  and  $B$ . Let  $t_1$  and  $t_2$  be the tangents to  $k_1$  and  $k_2$  at point  $A$ , and let  $t_1 \cap k_2 = \{A, C\}$ ,  $t_2 \cap k_1 = \{A, D\}$ . If  $E \in AB^{\rightarrow}$  such that  $AE = 2AB$ , prove that  $ACED$  is concyclic.
- (16) Let  $OL$  be the inner bisector of  $\angle POQ$ . A circle  $k$  passes through  $O$  and  $k \cap OP^{\rightarrow} = \{A\}$ ,  $k \cap OQ^{\rightarrow} = \{B\}$ ,  $k \cap OL^{\rightarrow} = \{C\}$ . Prove that, as  $k$  changes, the following ratio remains constant:

$$\frac{OA + OB}{OC}.$$

- (17) Let a circle  $k^*$  be inside a circle  $k$ ,  $k^* \cap k = \emptyset$ . We know that there exists a sequence of circles  $k_0, k_1, \dots, k_n$  such that  $k_i$  touches  $k, k^*$  and  $k_{i-1}$  for  $i = 1, 2, \dots, n + 1$  (here  $k_{n+1} = k_0$ .) Show that, instead of  $k_1$ , one can start with *any* circle  $k'_1$  tangent to both  $k$  and  $k^*$ , and still be able to fit a "ring" of  $n$  circles as above. What is  $n$  in terms of the radii of and the distance between the centers of  $k$  and  $k^*$ ?
- (18) Circles  $k_1, k_2, k_3$  touch pairwise, and all touch a line  $l$ . A fourth circle  $k$  touches  $k_1, k_2, k_3$ , so that  $k \cap l = \emptyset$ . Find the distance from the center of  $k$  to  $l$  given that radius of  $k$  is 1.