

Berkeley Math Circle Monthly Contest 7 – Solutions

1. Find all positive prime numbers p such that $p + 2$ and $p + 4$ are prime as well.

Hint. Show that for most prime numbers p , either $p + 2$ or $p + 4$ is divisible by 3.

Solution. For $p = 3$, $p + 2 = 5$, $p + 4 = 7$ and these are obviously prime. For $p > 3$, we know that p is not divisible by 3. The remainder of p when divided by 3 can be either 1 or 2. If it is one, then $p + 2$ is divisible by 3, if it is 2, then $p + 4$ is divisible by 3. Hence $p = 3$ is the only solution.

2. Let P be the point inside the square $ABCD$ such that $\triangle ABP$ is equilateral. Calculate the angle $\angle CPD$. Explain your answer!

Solution. The triangle DAP is isosceles because $AD = AP$ hence $\angle ADP = \angle APD = \frac{180^\circ - \angle DAP}{2} = 75^\circ$. Hence $\angle PDC = 15^\circ$. Similarly $\angle DCP = 15^\circ$ and hence $\angle CPD = 180^\circ - 2 \cdot 15^\circ = 150^\circ$.

3. Find at least one non-zero polynomial $P(x, y, z)$ such that $P(a, b, c) = 0$ for every three real numbers that satisfy $\sqrt[3]{a} + \sqrt[3]{b} = \sqrt[3]{c}$.

Remark. Polynomial in three variables refers to any expression built from x, y, z and numerals using only addition, subtraction, and multiplication. Parentheses or positive integer exponents, as in $x(y + z)^2$ are allowed since this can be expanded to $xyy + 2xyz + xzz$.

Solution. Cube both sides of the condition $\sqrt[3]{x} + \sqrt[3]{y} = \sqrt[3]{z}$:

$$x + 3\sqrt[3]{x}\sqrt[3]{x}\sqrt[3]{y} + 3\sqrt[3]{x}\sqrt[3]{y}\sqrt[3]{y} + y = z$$

$$3\sqrt[3]{x}\sqrt[3]{x}\sqrt[3]{y} + \sqrt[3]{x}\sqrt[3]{y}\sqrt[3]{y} = z - x - y$$

$$3\sqrt[3]{x}\sqrt[3]{y}(\sqrt[3]{x} + \sqrt[3]{y}) = z - x - y$$

But since $\sqrt[3]{x} + \sqrt[3]{y} = \sqrt[3]{z}$,

$$3\sqrt[3]{x}\sqrt[3]{y}\sqrt[3]{z} = z - x - y$$

$$27xyz = (z - x - y)^3.$$


Hence $P(x, y, z) = 27xyz - (z - x - y)^3$ is one such polynomial.

4. If $f(1) = 1$ and $f(1) + f(2) + \dots + f(n) = n^2 f(n)$ for every integer $n \geq 2$, evaluate $f(2008)$.

Solution. $n^2 f(n) - f(n) = f(1) + f(2) + \dots + f(n-1) = (n-1)^2 f(n-1)$ hence $f(n) = \frac{(n-1)^2}{n^2-1} f(n-1) = \frac{n-1}{n+1} f(n-1)$. Thus $f(2008) = \frac{2007}{2009} f(2007) = \frac{2007}{2009} \cdot \frac{2006}{2008} f(2006) = \frac{2007}{2009} \cdot \frac{2006}{2008} \cdot \frac{2005}{2007} f(2005) = \dots = \frac{2007!}{2009 \cdot 2008 \cdot \dots \cdot 4 \cdot 3} f(1) = \frac{2}{2009 \cdot 2008}$.

5. Given five vertices of a regular heptagon, construct the two remaining vertices using straightedge alone.

Solution. Let $A, B, C, D,$ and E be the known vertices and F and G the unknown vertices. The arrangement of $A, B, C, D,$ and E depends on the relative positions of F and G as shown in the diagram. The following construction applies to all three cases.



By connecting the intersections of BC with DE and BD with CE one finds l , the line of symmetry through G . Let AB and AC meet l in H and I respectively. By symmetry about l , HE and ID both pass through F . Now perform the same construction on B, C, D, E, F to get G .

Remark. In general, given five consecutive vertices A, B, C, D, E of a regular polygon or regular star polygon with at least seven sides, the next F can be found by this construction, except that l will not necessarily pass through G , although it is always the perpendicular bisector of CD, BE and AF and is a line of symmetry of the polygon.