Berkeley Math Circle Monthly Contest 7 Due April 1, 2008

Instructions

This contest consists of 5 problems, some of which are easier than the others. Every problem is worth 7 points. Please, write solution to every problem on a separate sheet of paper, and on top of each sheet include your name, grade and school, as well as the problem number and the contest number. Thus, the header on each sheet should look something like:

Solution to Problem 3 of BMC Monthly Contest 7 by Bart Simpson in grade 5 from Springfield Middle School, Springfield

If you submit more than one sheet for a specific problem, please, staple the sheets together to avoid getting them confused with someone else's solution. Please, do NOT staple together solutions to DIFFERENT problems, as they will be graded separately.

Carefully justify your answers to avoid losing points. Include all relevant explanations in words and all intermediate calculations. Answers without justification will receive no credit. However, good reasoning with minor calculational errors may receive a lot of points. Thus, submit solutions to as many problems as you can since partial credits will be awarded for sufficient progress on any particular problem.

Remember that you are NOT ALLOWED to consult or talk to anyone else about the problems, whether in person, on the phone, via e-mail, or other means of communication. You can consult any book that you wish. For more on the contest rules, please, check the BMC website at http://mathcircle.berkeley.edu.

Enjoy solving these problems and good luck!

Problems

1. Find all positive prime numbers p such that p + 2 and p + 4 are prime as well.

Hint. Show that for most prime numbers p, either p + 2 or p + 4 is divisible by 3.

- 2. Let P be the point inside the square ABCD such that $\triangle ABP$ is equilateral. Calculate the angle $\angle CPD$. Explain your answer!
- 3. Find at least one non-zero polynomial P(x, y, z) such that P(a, b, c) = 0 for every three real numbers that satisfy $\sqrt[3]{a} + \sqrt[3]{b} = \sqrt[3]{c}$.

Remark. Polynomial in three variables refers to any expression built from x, y, z and numerlas using only addition, subtraction, and multiplication. Parentheses or positive integer exponents, as in $x(y + z)^2$ are allowed since this can be expanded to xyy + 2xyz + xzz.

- 4. If f(1) = 1 and $f(1) + f(2) + \cdots + f(n) = n^2 f(n)$ for every integer $n \ge 2$, evaluate f(2008).
- 5. Given five vertices of a regular heptagon (7-gon), construct the two remaining vertices using straightedge alone.