

LINEAR RECURSIVE SEQUENCES

BJORN POONEN

1. SEQUENCES

A *sequence* is an infinite list of numbers, like

$$(1) \quad 1, 2, 4, 8, 16, 32, \dots$$

The numbers in the sequence are called its *terms*. The general form of a sequence is

$$a_1, a_2, a_3, \dots$$

where a_n is the n -th term of the sequence. In the example (1) above, $a_1 = 1$, $a_2 = 2$, $a_3 = 4$, and so on.

The notations $\{a_n\}$ or $\{a_n\}_{n=1}^{\infty}$ are abbreviations for

$$a_1, a_2, a_3, \dots$$

Occasionally the indexing of the terms will start with something other than 1. For example, $\{a_n\}_{n=0}^{\infty}$ would mean

$$a_0, a_1, a_2, \dots$$

(In this case a_n would be the $(n + 1)$ -st term.)

For some sequences, it is possible to give an *explicit formula* for a_n : this means that a_n is expressed as a function of n . For instance, the sequence (1) above can be described by the explicit formula $a_n = 2^{n-1}$.

2. RECURSIVE DEFINITIONS

An alternative way to describe a sequence is to list a few terms and to give a rule for computing the rest of the sequence. Our example (1) above can be described by the starting value $a_1 = 1$ and the rule $a_{n+1} = 2a_n$ for integers $n \geq 1$. Starting from $a_1 = 1$, the rule implies that

$$\begin{aligned} a_2 &= 2a_1 = 2(1) = 2 \\ a_3 &= 2a_2 = 2(2) = 4 \\ a_4 &= 2a_3 = 2(4) = 8, \end{aligned}$$

and so on; each term in the sequence can be computed recursively in terms of the terms previously computed. A rule such as this giving the next term in terms of earlier terms is also called a *recurrence relation* (or simply *recurrence*).

3. LINEAR RECURSIVE SEQUENCES

A sequence $\{a_n\}$ is said to satisfy the *linear recurrence* with coefficients c_k, c_{k-1}, \dots, c_0 if

$$(2) \quad c_k a_{n+k} + c_{k-1} a_{n+k-1} + \dots + c_1 a_{n+1} + c_0 a_n = 0$$

holds for all integers n for which this makes sense. (If the sequence starts with a_1 , then this means for $n \geq 1$.) The integer k is called the *order* of the linear recurrence.

A *linear recursive sequence* is a sequence of numbers a_1, a_2, a_3, \dots satisfying some linear recurrence as above with $c_k \neq 0$ and $c_0 \neq 0$. For example, the sequence (1) satisfies

$$a_{n+1} - 2a_n = 0$$

for all integers $n \geq 1$, so it is a linear recursive sequence satisfying a recurrence of order 1, with $c_1 = 1$ and $c_0 = -2$.

Requiring $c_k \neq 0$ guarantees that the linear recurrence can be used to express a_{n+k} as a linear combination of earlier terms:

$$a_{n+k} = -\frac{c_{k-1}}{c_k} a_{n+k-1} - \dots - \frac{c_1}{c_k} a_{n+1} - \frac{c_0}{c_k} a_n.$$

The requirement $c_0 \neq 0$ lets one express a_n as a linear combination of *later* terms:

$$a_n = -\frac{c_k}{c_0} a_{n+k} - \frac{c_{k-1}}{c_0} a_{n+k-1} - \dots - \frac{c_1}{c_0} a_{n+1}.$$

This lets one *define* a_0, a_{-1} , and so on, to obtain a *doubly infinite sequence*

$$\dots, a_{-2}, a_{-1}, a_0, a_1, a_2, \dots$$

that now satisfies the same linear recurrence for *all* integers n , positive or negative.

4. CHARACTERISTIC POLYNOMIALS

The *characteristic polynomial* of a linear recurrence

$$c_k a_{n+k} + c_{k-1} a_{n+k-1} + \dots + c_1 a_{n+1} + c_0 a_n = 0$$

is defined to be the polynomial

$$c_k x^k + c_{k-1} x^{k-1} + \dots + c_1 x + c_0.$$

For example, the characteristic polynomial of the recurrence $a_{n+1} - 2a_n = 0$ satisfied by the sequence (1) is $x - 2$.

Here is another example: the famous *Fibonacci sequence*

$$\{F_n\}_{n=0}^\infty = 0, 1, 1, 2, 3, 5, 8, 13, \dots$$

which can be described by the starting values $F_0 = 0, F_1 = 1$ and the recurrence relation

$$(3) \quad F_n = F_{n-1} + F_{n-2} \quad \text{for all } n \geq 2.$$

To find the characteristic polynomial, we first need to rewrite the recurrence relation in the form (2). The relation (3) is equivalent to

$$(4) \quad F_{n+2} = F_{n+1} + F_n \quad \text{for all } n \geq 0.$$

Rewriting it as

$$(5) \quad F_{n+2} - F_{n+1} - F_n = 0$$

shows that $\{F_n\}$ is a linear recursive sequence satisfying a recurrence of order 2, with $c_2 = 1$, $c_1 = -1$, and $c_0 = -1$. The characteristic polynomial is $x^2 - x - 1$.

5. IDEALS AND MINIMAL CHARACTERISTIC POLYNOMIALS

The same sequence can satisfy many different linear recurrences. For example, doubling (5) shows the Fibonacci sequence also satisfies

$$2F_{n+2} - 2F_{n+1} - 2F_n = 0,$$

which is a linear recurrence with characteristic polynomial $2x^2 - 2x - 2$. It also satisfies

$$F_{n+3} - F_{n+2} - F_{n+1} = 0,$$

and adding these two relations, we find that $\{F_n\}$ also satisfies

$$F_{n+3} + F_{n+2} - 3F_{n+1} - 2F_n = 0$$

which has characteristic polynomial $x^3 + x^2 - 3x - 2 = (x + 2)(x^2 - x - 1)$.

Now consider an arbitrary sequence $\{a_n\}$. Let I be the set of characteristic polynomials of *all* linear recurrences satisfied by $\{a_n\}$. Then

- (a) If $f(x) \in I$ and $g(x) \in I$ then $f(x) + g(x) \in I$.
- (b) If $f(x) \in I$ and $h(x)$ is any polynomial, then $h(x)f(x) \in I$.

In general, a nonempty set I of polynomials satisfying (a) and (b) is called an *ideal*.

Fact from algebra: Let I be an ideal of polynomials. Then either $I = \{0\}$ or else there is a unique monic polynomial $f(x) \in I$ such that

$$I = \text{the set of polynomial multiples of } f(x) = \{h(x)f(x) \mid h(x) \text{ is a polynomial}\}.$$

(A polynomial is *monic* if the coefficient of the highest power of x is 1.)

This fact, applied to the ideal of characteristic polynomials of a linear recursive sequence $\{a_n\}$ shows that there is always a *minimal characteristic polynomial* $f(x)$, which is the monic polynomial of lowest degree in I . It is the characteristic polynomial of the lowest order nontrivial linear recurrence satisfied by $\{a_n\}$. The characteristic polynomial of any other linear recurrence satisfied by $\{a_n\}$ is a polynomial multiple of $f(x)$.

The *order* of a linear recursive sequence $\{a_n\}$ is defined to be the lowest order among all (nontrivial) linear recurrences satisfied by $\{a_n\}$. The order also equals the degree of the minimal characteristic polynomial. For example, as we showed above, $\{F_n\}$ satisfies

$$F_{n+3} + F_{n+2} - 3F_{n+1} - 2F_n = 0,$$

but we also know that

$$F_{n+2} - F_{n+1} - F_n = 0,$$

and it is easy to show that $\{F_n\}$ cannot satisfy a linear recurrence of order less than 2, so $\{F_n\}$ is a linear recursive sequence of order 2, with minimal characteristic polynomial $x^2 - x - 1$.

6. THE MAIN THEOREM

Theorem 1. Let $f(x) = c_k x^k + \dots + c_0$ be a polynomial with $c_k \neq 0$ and $c_0 \neq 0$. Factor $f(x)$ over the complex numbers as

$$f(x) = c_k(x - r_1)^{m_1}(x - r_2)^{m_2} \dots (x - r_\ell)^{m_\ell},$$

where r_1, r_2, \dots, r_ℓ are distinct nonzero complex numbers, and m_1, m_2, \dots, m_ℓ are positive integers. Then a sequence $\{a_n\}$ satisfies the linear recurrence with characteristic polynomial $f(x)$ if and only if there exist polynomials $g_1(n), g_2(n), \dots, g_\ell(n)$ with $\deg g_i \leq m_i - 1$ for $i = 1, 2, \dots, \ell$ such that

$$a_n = g_1(n)r_1^n + \dots + g_\ell(n)r_\ell^n \quad \text{for all } n.$$

Here is an important special case.

Corollary 2. *Suppose in addition that $f(x)$ has no repeated factors; in other words suppose that $m_1 = m_2 = \dots = m_\ell = 1$. Then $f(x) = c_k(x - r_1)(x - r_2) \dots (x - r_\ell)$ where r_1, r_2, \dots, r_ℓ are distinct nonzero complex numbers (the roots of f). Then $\{a_n\}$ satisfies the linear recurrence with characteristic polynomial $f(x)$ if and only if there exist constants B_1, B_2, \dots, B_ℓ (not depending on n) such that*

$$a_n = B_1 r_1^n + \dots + B_\ell r_\ell^n \quad \text{for all } n.$$

7. EXAMPLE: SOLVING A LINEAR RECURRENCE

Suppose we want to find an explicit formula for the sequence $\{a_n\}$ satisfying $a_0 = 1$, $a_1 = 4$, and

$$(6) \quad a_{n+2} = \frac{a_{n+1} + a_n}{2} \text{ for } n \geq 0.$$

Since $\{a_n\}$ satisfies a linear recurrence with characteristic polynomial $x^2 - \frac{1}{2}x - \frac{1}{2} = (x - 1)(x + \frac{1}{2})$, we know that there exist constants A and B such that

$$(7) \quad a_n = A(1)^n + B\left(-\frac{1}{2}\right)^n$$

for all n . The formula (7) is called the *general solution* to the linear recurrence (6). To find the *particular solution* with the correct values of A and B , we use the known values of a_0 and a_1 :

$$\begin{aligned} 1 = a_0 &= A(1)^0 + B\left(-\frac{1}{2}\right)^0 = A + B \\ 4 = a_1 &= A(1)^1 + B\left(-\frac{1}{2}\right)^1 = A - B/2. \end{aligned}$$

Solving this system of equations yields $A = 3$ and $B = -2$. Thus the particular solution is

$$a_n = 3 - 2\left(-\frac{1}{2}\right)^n.$$

(As a check, one can try plugging in $n = 0$ or $n = 1$.)

8. EXAMPLE: THE FORMULA FOR THE FIBONACCI SEQUENCE

As we worked out earlier, $\{F_n\}$ satisfies a linear recurrence with characteristic polynomial $x^2 - x - 1$. By the quadratic formula, this factors as $(x - \alpha)(x - \beta)$ where $\alpha = (1 + \sqrt{5})/2$ is the golden ratio, and $\beta = (1 - \sqrt{5})/2$. The main theorem implies that there are constants A and B such that

$$F_n = A\alpha^n + B\beta^n$$

for all n . Using $F_0 = 0$ and $F_1 = 1$ we obtain

$$0 = A + B, \quad 1 = A\alpha + B\beta.$$

Solving for A and B yields $A = 1/(\alpha - \beta)$ and $B = -1/(\alpha - \beta)$, so

$$F_n = \frac{\alpha^n - \beta^n}{\alpha - \beta} = \frac{1}{\sqrt{5}} \left[\left(\frac{1 + \sqrt{5}}{2} \right)^n - \left(\frac{1 - \sqrt{5}}{2} \right)^n \right]$$

for all n .

9. EXAMPLE: FINDING A LINEAR RECURRENCE FROM AN EXPLICIT FORMULA

Let $a_n = (n + 2^n)F_n$, where $\{F_n\}$ is the Fibonacci sequence. Then by the explicit formula for F_n ,

$$\begin{aligned} a_n &= (n + 2^n) \left(\frac{\alpha^n - \beta^n}{\alpha - \beta} \right) \\ &= \left[\left(\frac{1}{\alpha - \beta} \right) n \right] \alpha^n + \left[\left(\frac{-1}{\alpha - \beta} \right) n \right] \beta^n + \left(\frac{1}{\alpha - \beta} \right) (2\alpha)^n + \left(\frac{-1}{\alpha - \beta} \right) (2\beta)^n. \end{aligned}$$

By Theorem 1, $\{a_n\}$ satisfies a linear recurrence with characteristic polynomial

$$\begin{aligned} (x - \alpha)^2(x - \beta)^2(x - 2\alpha)(x - 2\beta) &= (x^2 - x - 1)^2 [x^2 - 2(\alpha + \beta) + 4\alpha\beta] \\ &= (x^2 - x - 1)^2(x^2 - 2x - 4) \\ &= x^6 - 4x^5 - x^4 + 12x^3 + x^2 - 10x + 4, \end{aligned}$$

where we have used the identity $x^2 - (\alpha + \beta)x + \alpha\beta = x^2 - x - 1$ to compute $\alpha + \beta$ and $\alpha\beta$. In other words,

$$a_{n+6} - 4a_{n+5} - a_{n+4} + 12a_{n+3} + a_{n+2} - 10a_{n+1} + 4a_n = 0$$

for all n . In fact, we have found the minimal characteristic polynomial, since if the actual minimal characteristic polynomial were a proper divisor of $(x^2 - x - 1)^2(x^2 - 2x - 4)$, then according to Theorem 1, the explicit formula for a_n would have had a different, simpler form.

10. INHOMOGENEOUS RECURRENCE RELATIONS

Suppose we wanted an explicit formula for a sequence $\{a_n\}$ satisfying $a_0 = 0$, and

$$(8) \quad a_{n+1} - 2a_n = F_n \quad \text{for } n \geq 0,$$

where $\{F_n\}$ is the Fibonacci sequence as usual. This is not a linear recurrence in the sense we have been talking about (because of the F_n on the right hand side instead of 0), so our usual method does not work. A recurrence of this type, linear except for a function of n on the right hand side, is called an *inhomogeneous recurrence*.

We can solve inhomogeneous recurrences explicitly when the right hand side is itself a linear recursive sequence. In our example, $\{a_n\}$ also satisfies

$$(9) \quad a_{n+2} - 2a_{n+1} = F_{n+1}$$

and

$$(10) \quad a_{n+3} - 2a_{n+2} = F_{n+2}.$$

Subtracting (8) and (9) from (10) yields

$$a_{n+3} - 3a_{n+2} + a_{n+1} + 2a_n = F_{n+2} - F_{n+1} - F_n = 0.$$

Thus $\{a_n\}$ is a linear recursive sequence after all! The characteristic polynomial of this new linear recurrence is $x^3 - 3x^2 + x + 2 = (x - 2)(x^2 - x - 1)$, so by Theorem 1, there exist constants A, B, C such that

$$a_n = A \cdot 2^n + B\alpha^n + C\beta^n$$

for all n . Now we can use $a_0 = 0$, and the values $a_1 = 0$ and $a_2 = 1$ obtained from (8) to determine A, B, C . After some work, one finds $A = 1$, $B = -\alpha^2/(\alpha - \beta)$, and $C = \beta^2/(\alpha - \beta)$, so $a_n = 2^n - F_{n+2}$.

If $\{x_n\}$ is any other sequence satisfying

$$(11) \quad x_{n+1} - 2x_n = F_n$$

but not necessarily $x_0 = 0$, then subtracting (8) from (11) shows that the sequence $\{y_n\}$ defined by $y_n = x_n - a_n$ satisfies $y_{n+1} - 2y_n = 0$ for all n , so $y_n = D \cdot 2^n$ for some number D . Hence the *general solution* of (11) has the form

$$x_n = 2^n - F_{n+2} + D \cdot 2^n,$$

or more simply,

$$x_n = E \cdot 2^n - F_{n+2},$$

where E is some constant.

In general, this sort of argument proves the following.

Theorem 3. *Let $\{b_n\}$ be a linear recursive sequence satisfying a recurrence with characteristic polynomial $f(x)$. Let $g(x) = c_k x^k + c_{k-1} x^{k-1} + \cdots + c_1 x + c_0$ be a polynomial. Then every solution $\{x_n\}$ to the inhomogeneous recurrence*

$$(12) \quad c_k x_{n+k} + c_{k-1} x_{n+k-1} + \cdots + c_1 x_{n+1} + c_0 x_n = b_n$$

also satisfies a linear recurrence with characteristic polynomial $f(x)g(x)$. Moreover, if $\{x_n\} = \{a_n\}$ is one particular solution to (12), then all solutions have the form $x_n = a_n + y_n$, where $\{y_n\}$ ranges over the solutions of the linear recurrence

$$c_k y_{n+k} + c_{k-1} y_{n+k-1} + \cdots + c_1 y_{n+1} + c_0 y_n = 0.$$

11. THE MAHLER-LECH THEOREM

Here is a deep theorem about linear recursive sequences:

Theorem 4 (Mahler-Lech theorem). *Let $\{a_n\}$ be a linear recursive sequence of complex numbers, and let c be a complex number. Then there exists a finite (possibly empty) list of arithmetic progressions T_1, T_2, \dots, T_m and a finite (possibly empty) set S of integers such that*

$$\{n \mid a_n = c\} = S \cup T_1 \cup T_2 \cup \cdots \cup T_m.$$

Warning: don't try to prove this at home! This is *very* hard to prove. The proof uses “ p -adic numbers.”

12. PROBLEMS

There are a lot of problems here. Just do the ones that interest you.

- (1) If the Fibonacci sequence is extended to a doubly infinite sequence satisfying the same linear recurrence, then what will F_{-4} be? (Is it easier to do this using the recurrence, or using the explicit formula?)
- (2) Find the smallest degree polynomial that could be the minimal characteristic polynomial of a sequence that begins

$$2, 5, 18, 67, 250, 933, \dots$$

Assuming that the sequence *is* a linear recursive sequence with this characteristic polynomial, find an explicit formula for the n -th term.

- (3) Suppose that $a_n = n^2 + 3n + 7$ for $n \geq 1$. Prove that $\{a_n\}$ is a linear recursive sequence, and find its minimal characteristic polynomial.
- (4) Suppose $a_1 = a_2 = a_3 = 1$, $a_4 = 3$, and $a_{n+4} = 3a_{n+2} - 2a_n$ for $n \geq 1$. Prove that $a_n = 1$ if and only if n is odd or $n = 2$. (This is an instance of the Mahler-Lech theorem: for this sequence, one would take $S = \{2\}$ and $T_1 = \{1, 3, 5, 7, \dots\}$.)
- (5) Suppose $a_0 = 2$, $a_1 = 5$, and $a_{n+2} = (a_{n+1})^2(a_n)^3$ for $n \geq 0$. (This is a recurrence relation, but not a linear recurrence relation.) Find an explicit formula for a_n .
- (6) Suppose $\{a_n\}$ is a sequence such that $a_{n+2} = a_{n+1} - a_n$ for all $n \geq 1$. Given that $a_{38} = 7$ and $a_{55} = 3$, find a_1 . (Hint: it is possible to solve this problem with very little calculation.)
- (7) Let θ be a fixed real number, and let $a_n = \cos(n\theta)$ for integers $n \geq 1$. Prove that $\{a_n\}$ is a linear recursive sequence, and find the minimal characteristic polynomial. (Hint: if you know the definition of $\cos x$ in terms of complex exponentials, use that. Otherwise, use the sum-to-product rule for the sum of cosines $\cos(n\theta) + \cos((n+2)\theta)$. For most but not all θ , the degree of the minimal characteristic polynomial will be 2.)
- (8) Give an example of a sequence that is *not* a linear recursive sequence, and prove that it is not one.
- (9) Given a finite set S of positive integers, show that there exists a linear recursive sequence

$$a_1, a_2, a_3, \dots$$

such that $\{n \mid a_n = 0\} = S$.

- (10) A student tosses a fair coin and scores one point for each head that turns up, and two points for each tail. Prove that the probability of the student scoring n points at some time in a sequence of n tosses is $\frac{1}{3} \left(2 + \left(-\frac{1}{2}\right)^n\right)$.
- (11) Let F_n denote the n -th Fibonacci number. Let $a_n = (F_n)^2$. Prove that a_1, a_2, a_3, \dots is a linear recursive sequence, and find its minimal characteristic polynomial.
- (12) Prove the “fact from algebra” mentioned above in Section 5. (Hint: if $I \neq \{0\}$, pick a nonzero polynomial in I of smallest degree, and multiply it by a constant to get a monic polynomial $f(x)$. Use long division of polynomials to show that anything else in I is a polynomial multiple of $f(x)$.)
- (13) Suppose that a_1, a_2, \dots is a linear recursive sequence. For $n \geq 1$, let $s_n = a_1 + a_2 + \dots + a_n$. Prove that $\{s_n\}$ is a linear recursive sequence.
- (14) Suppose $\{a_n\}$ and $\{b_n\}$ are linear recursive sequences. Let $c_n = a_n + b_n$ and $d_n = a_n b_n$ for $n \geq 1$.
 - (a) Prove that $\{c_n\}$ and $\{d_n\}$ also are linear recursive sequences.

(b) Suppose that the minimal characteristic polynomials for $\{a_n\}$ and $\{b_n\}$ are $x^2 - x - 2$ and $x^2 - 5x + 6$, respectively. What are the possibilities for the minimal characteristic polynomials of $\{c_n\}$ and $\{d_n\}$?

- (15) Suppose that $\{a_n\}$ and $\{b_n\}$ are linear recursive sequences. Prove that

$$a_1, b_1, a_2, b_2, a_3, b_3, \dots$$

also is a linear recursive sequence.

- (16) Use the Mahler-Lech theorem to prove the following generalization. Let $\{a_n\}$ be a linear recursive sequence of complex numbers, and let $p(x)$ be a polynomial. Then there exists a finite (possibly empty) list of arithmetic progressions T_1, T_2, \dots, T_m and a finite (possibly empty) set S of integers such that

$$\{n \mid a_n = p(n)\} = S \cup T_1 \cup T_2 \cup \dots \cup T_m.$$

(Hint: let $b_n = a_n - p(n)$.)

- (17) (1973 USAMO, no. 2) Let $\{X_n\}$ and $\{Y_n\}$ denote two sequences of integers defined as follows:

$$X_0 = 1, X_1 = 1, X_{n+1} = X_n + 2X_{n-1} \quad (n = 1, 2, 3, \dots),$$

$$Y_0 = 1, Y_1 = 7, Y_{n+1} = 2Y_n + 3Y_{n-1} \quad (n = 1, 2, 3, \dots).$$

Thus, the first few terms of the sequences are:

$$X : 1, 1, 3, 5, 11, 21, \dots,$$

$$Y : 1, 7, 17, 55, 161, 487, \dots$$

Prove that, except for the “1,” there is no term which occurs in both sequences.

- (18) (1963 IMO, no. 4) Find all solutions x_1, x_2, x_3, x_4, x_5 to the system

$$x_5 + x_2 = yx_1$$

$$x_1 + x_3 = yx_2$$

$$x_2 + x_4 = yx_3$$

$$x_3 + x_5 = yx_4$$

$$x_4 + x_1 = yx_5,$$

where y is a parameter. (Hint: define $x_6 = x_1, x_7 = x_2$, etc., and find two different linear recurrences satisfied by $\{x_n\}$.)

- (19) (1967 IMO, no. 6) In a sports contest, there were m medals awarded on n successive days ($n > 1$). On the first day, one medal and $1/7$ of the remaining $m - 1$ medals were awarded. On the second day, two medals and $1/7$ of the now remaining medals were awarded; and so on. On the n -th and last day, the remaining n medals were awarded. How many days did the contest last, and how many medals were awarded altogether?

- (20) (1974 IMO, no. 3) Prove that the number $\sum_{k=0}^n \binom{2n+1}{k+1} 2^{3k}$ is not divisible by 5 for any integer $n \geq 0$.

- (21) (1980 USAMO, no. 3) Let $F_r = x^r \sin(rA) + y^r \sin(rB) + z^r \sin(rC)$, where x, y, z, A, B, C are real and $A + B + C$ is an integral multiple of π . Prove that if $F_1 = F_2 = 0$, then $F_r = 0$ for all positive integral r .

©Berkeley Math Circle

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720-3840, USA

E-mail address: `poonen@math.berkeley.edu`